

DECIPHERING THE MICROBIAL DIVERSITY OF TATTAPANI HOT WATER SPRING USING METAGENOMIC APPROACH

MAHAJAN MAHESH MOHANRAO¹, DUSHYANT PRATAP SINGH²,
KUMAR KANIKA³, ETIKA GOYAL⁴ & AMIT K. SINGH⁵

^{1,4,5}Research Scholar, NRCPB, New Delhi, India

²Research Scholar, JJT University, Rajasthan, India

³Senior Scientist, NRCPB, New Delhi, India

ABSTRACT

Heat stress due to changing environmental conditions is a problem as it negatively affects plant growth and development leading to reduction in crop yield. Thermal springs are major source of potentially significant but untapped microbial diversity encoding novel genes related to high temperature tolerance. The genes involved in the heat tolerance mechanism of thermophiles can serve as a valuable resource to improve high temperature tolerance in crop plants. Metagenomics bypasses the need for isolation or cultivation of microorganisms as it deals with direct isolation of nucleic acids from environmental samples. The present study was designed to analyse the profile of microbial community inhabiting hot water spring Tattapani, Himachal Pradesh, India. The spring water temperature at the discharge point was recorded as 65°C and pH 7.5. Two sets of universal 16S rRNA gene specific primers namely U789F-U1053R and MGUF-MGUR were used to amplify partial 16S rRNA gene (from V5-V6 and V3-V6 regions respectively) from metagenomic DNA. Sequencing and bioinformatic analysis revealed that bacterial phylum Proteobacteria was most abundant, constituting 36% and 40% of the total bacteria identified using V5-V6 and V3-V6 regions, respectively. Other phyla Thermodesulfobacteria (9.3%), Firmicutes (7.5%), Deinococcus-thermus (7%), Bacteroidates (2.5%), Aquificae (2%) were assigned to sequences amplified from V5-V6 region while phyla Firmicutes (18%), Bacteroidates (6%), Chloroflexi (2%) were assigned to sequences amplified from V3-V6 region. A significant number of sequences (30.5% and 20.5%) from region V3-V6 and V5-V6 respectively did not show homology with existing database and were recorded as unclassified Bacteria. The information generated in this study will help us in understanding the microbial ecology of hot springs.

KEYWORDS: Microbial Diversity, Metagenomics, Hot Water Spring

Received: Mar 21, 2016; **Accepted:** Apr 06, 2016; **Published:** Apr 08, 2016; **Paper Id.:** IJASRAPR201650

INTRODUCTION

Natural environments are shelter for large diversity of microorganisms; most of these microorganisms are unexplored and belong to untapped genetic resource (Nichols, 2002). Extreme environments such as hot water springs, arctic regions, highly saline or alkaline regions, deep oceans etc. are a major source of potentially significant but untapped microbial diversity. Study of microbial diversity in such extreme environments is necessary because, different physico-chemical conditions, biogeography and geological history, have different microbial phenotypes (Narayan, 2008; Liu et al., 2009). Of the total microorganisms present in environment only 0.1 to 1% can be cultured by conventional laboratory methods (Amann et al., 1995). Thus remaining ~99% of microbes are unculturable. To overcome this problem culture independent approach (metagenomics) is used that

enable us to investigate microbial diversity from diverse habitats. Metagenomic approach based on 16S rRNA gene analysis through its amplification and sequencing provides information about microbial community present in an environment.

The partial 16S rRNA gene sequence based metagenomic approach has been extensively used to study microbiota present in hot water springs. These includes, Sungai klah hot spring, Malaysia (chan et al., 2015), Tuwa hot spring, India (Mangrola et al., 2015) hot spring in Tibetan plateau, China (Huang et al., 2011), hot spring in Tengchong County in Yunnan Province, China (Briggs et al., 2013; Song et al., 2009, 2010; Jiang et al., 2010), different geothermal hot springs in Tunisia (Sayeh et al., 2010) Long Valley Caldera near Mammoth Lakes, CA, USA (Vick et al., 2010), hot springs in Seltun and Hveradalir geothermal areas, Iceland (Aguilera et al., 2010; Reigstad et al., 2009; Marteinson et al., 2001), acidic hot spring, Kawah Hujan B, Indonesia (Aditiawati et al., 2009), hot springs located at Yellowstone National Park, USA (Meyer-Dombard et al., 2005; Hall et al., 2008; Mitchell 2009), Soldhar and Ringigad hot springs, Uttaranchal, India (Kumar et al., 2004) and Kamchatka hot spring, Russia (Bonch-Osmolovskaya et al., 1999; Reigstad et al., 2009). These studies revealed that temperature, pH and biogeography are the important factors affecting the microbial diversity of the hot water springs (Huang et al., 2011).

Though large number of studies has been carried out on hot springs habitats, there is a lack of information on microbial communities inhabiting hot water springs in India. In this backdrop, present study was planned to study the microbial diversity inhabiting the hot water spring at Tattapani, Himachal Pradesh, India. The Tattapani hot water spring is located on the bank of river Satluj. Metagenomic DNA from the hot spring water was isolated and partial 16S rRNA gene was amplified using primers designed from V3-6 and V5-6 hypervariable regions of prokaryotic 16S rRNA genes. PCR amplified partial 16S rRNA gene fragments were cloned and sequenced. Different bioinformatic tools were used to investigate the microbial community structure of Tattapani hot water spring.

MATERIALS AND METHODS

Study Area and Sampling

The Tattapani hot water spring is located at an altitude of 655 meters above sea level, on the bank of river Satluj, Mandi district, Himachal Pradesh, India, having geographical coordinates 31° 14' 56"N, 77° 5' 10"E (Figure 1a and 1b). The location of hot spring is geographically very unique as the temperature of river water was recorded near about 5°C-6°C while the temperature of hot water of the reservoir was ~65°C. Hot water samples for microbial diversity analysis were collected from the hot springs oozing out from different places on the banks of river. Water samples were pooled and used for metagenomic DNA extraction. White and light yellow coloured salt and sulphur deposits were observed all around the surrounding stones of the hot spring along with some algae which formed mats along hot springs. Different physico-chemical water quality parameters, namely; temperature, pH, electrical conductivity (EC) and total dissolved solid (TDS) were measured on-site using the relevant field meters.

Metagenomic DNA Extraction

Pooled hot spring water sample were filtered using Millipore filter membrane having pore size 0.22µm through vacuum filtration system at collection site. The microbial mass was concentrated on the surface of the filter papers which were further used to isolate the metagenomic DNA. The filter papers were sliced in to smaller pieces, placed in aseptic centrifuge tubes and processed for extraction of metagenomic DNA using RapidWater™ DNA Isolation (MO BIO Laboratories, Inc.) as per the manufacturer's protocol. DNA extraction was performed multiple times, extracted DNA was

pooled to get concentrated DNA. The quality and quantity of DNA was evaluated by using agarose gel electrophoresis and Nanodrop™ 1000 spectrophotometer (Thermo Scientific, Wilmington, DE, USA).

Targeted 16S rRNA Clone Library Preparation

The PCR was performed using GOTaq® green master mix (Promega, USA) for amplification of partial 16S rRNA gene using diluted metagenomic DNA as template in Eppendorf master cycler. The oligonucleotide primer sets used in this study are shown in Table 1. Thermal cycler conditions used for primer set MGUF/MGUR were as follows: 4 min. of initial denaturation at 94°C; followed by 35 amplification cycles of 1 min. at 94°C, 45 sec. at 60.7°C, 1 min. at 72°C; with a final 10 min of prolonged extension at 72°C. The conditions for primer set U789F/U1053R were similar to the above conditions except for annealing and extension which were 58.6°C for 45 sec., 72°C for 40 sec. respectively. Amplified fragments were visualized on 1.2% agarose gel. The amplicons were gel purified using the column based MinElute® Gel Extraction Kit (Qiagen, Germany) as per the manufacturers protocol.

Table 1: The Primer Sequence Used for Amplification of Partial 16S rRNA Gene

Primer Name	Primer Sequence	Expected Size of Amplicons	V Region	Reference
MGUF MGUR	5' TGCCAGCAGCCGCGGTA 3' 5' GACGGGCGGTGTGTACAA 3'	850bp	V3-V6	Wang, Y. and Qian, P. (2009)
U789F U1053R	5' TAGATACCCSSGTAGTTC 3' 5' CTGACGRCRGCCATGC 3'	270bp	V5-V6	Wang, Y. and Qian, P. (2009)

Cloning and Transformation of Partial 16S rRNA Gene Fragment

Gel purified PCR products were ligated with pGEMT easy vector (Promega, USA). The reaction was incubated at 4°C overnight as recommended in manufactures protocol. Electrocompetent DH5α *E. coli* cells were transformed with ligated product using standard electroporation protocol (Sambrook and Russell, 2001). The transformed cells were picked using blue/white screening. The transformants were confirmed for the presence of the insert using colony PCR. Positive clones were used for isolation of plasmid DNA using R.E.A.L® prep 96 Plasmid Isolation Kit (Qiagen, Germany)) as per the manufacturer's protocol.

in silico Analysis of 16S rRNA Gene Sequences

The partial 16S rRNA clones were sequenced and data were classified using the RDP Naive Bayesian rRNA Classifier Version 2.10, October 2014, RDP training set 14, based on nomenclatural taxonomy and Bergey's Manual with an 80% confidence threshold on the RDP-database. DECIPHER, a search based approach was used to identify the presence of possible chimeric sequences from all 16S rRNA gene sequences (Wright et al. 2012). Bioinformatics analysis of the sequences was carried out to identify homology match. After excluding redundant and false-positive sequences, nucleic acid homology searches were performed against nucleotide databases at the National Center for Biotechnology Information (NCBI) using the Basic Local Alignment Search Tool (BLASTN) <http://www.ncbi.nlm.nih.gov/BLAST> to provide gene annotation. Phylogenetic trees of selected sequences were constructed from distance matrices by the neighbor-joining method (Naruya et al., 1987), using MEGA6 tool (Figure 3a and 3b). The node reproducibility for tree topology was estimated by bootstrap analysis, which included 1000 replicate data sets.

Nucleotide Sequence Accession Numbers

The nucleotide sequences from V3-V6 region and V5-V6 region have been deposited in the GenBank (NCBI)

database under temporary accession numbers KU970288 - KU970336 and KU970337 - KU970418 respectively.

RESULTS AND DISCUSSIONS

Physico-Chemical Analysis of Spring Water Sample

The physico-chemical parameters of the spring water such as temperature and pH are important factors that determine microbial diversity in hot springs (Hou et al., 2013; Cole et al., 2013). High temperature of water exerts stress on microbial species; therefore only tolerant species can survive (Abou-Shanab et al., 2007). In present study water sampling was done at different spots of hot springs present on the river bed and samples were pooled to obtain final representative water sample. The spring water was clear in appearance with smell of sulphur. The pH of water was 7.5 which showed that it was slightly alkaline in nature. The Physico-chemical parameters of water are as summarized Table 2

Table 2: Physico-Chemical Parameters of Tattapani Hot Water Spring

Physico-Chemical Parameters	Observations
Temperature	65°C
pH	7.5
EC	17.3 00 dS/m
TDS	8400 mg/L
Odour	Rotten egg (qualitative)
Colour	Normal (Hazen scale unit)
Appearance	Clear (qualitative)
Taste	Salty (qualitative)
Sulphate	112 mg/L

Extraction of Metagenome and Cloning of Partial 16S rRNA Gene

Number of protocols has been standardized for extraction of metagenomic DNA (Zhou et al., 1996; Sharma et al., 2007; Aditiawati et al., 2009) and there are various kits available for extraction and purification of DNA from environmental water samples. Some of them are easy to use and reproducible. However, none of these methods are universally applicable to all water type. Each water type requires optimization of DNA extraction protocol. In the present study metagenomic DNA isolation was done by Rapid™ DNA Isolation (Mo Bio Laboratories', Inc.) (Figure 2a). The isolated DNA was of good quality with no visible shearing and its concentration was calculated as ~90 ng/μl. The purity ratio was measured as 1.87 at wavelength 260/280 using Nanodrop™ 1000 spectrophotometer (Thermo Scientific, Wilmington, DE, USA). Purified metagenomic DNA was used for PCR amplification of partial 16S rRNA gene from two variable regions (Figure 2b). Amplified 16S rRNA gene fragments were cloned in pGEM®-T easy cloning vector. After blue white screen around 105 positive clones were picked having 850bp 16S rRNA gene fragment and 161 clones were picked having 270bp 16S rRNA gene fragment.

Taxonomic Profile Analysis Using V5-6 and V3-6 Region of 16S rRNA

To investigate taxonomic profile present in environmental samples, 16S rRNA gene analysis with its highly conserved and variable sequence provides a major target region for studying microbial diversity (Muyzer et al., 1993; Becker et al., 2004; Wang and Qian, 2009). In recent metagenomic studies, the number of phylotypes in the same number of 16S rRNA gene sequences varied substantially for samples from different environments and geographical sites (Sogin, 2006; Huber, 2007; Roesch, 2007). Primer usage is undoubtedly one of the most critical limiting factors affecting 16S rRNA gene analysis (Armougom and Raoult, 2009). The amplification efficacy of the different 16S rRNA gene primers used can lead to under or over estimation of some microbial phyla. In various studies V3 and V6 have been most

extensively examined regions (Huse et al., 2008; Wang et al., 2009). However different primers targeting different regions of 16S rRNA gene were used for amplification in different studies. In present study, two sets of primers targeting V3-V6 region were used for 16S rRNA gene amplification (Wang et al., 2009). To study microbial diversity of Kawah Hujan B hot spring, Indonesia, primers were designed from variable region V7-V8 (Aditiawati et al., 2009), for Malaysian hot spring amplicons of V3-V4 region were amplified (Chan et al., 2015), regions V1-3 and V4-7 were amplified for bacterial diversity analysis of Siloam hot water spring (Tekere et al., 2011). Different sets of primers lead to difference in the capture depth of the microbes in different environmental samples because of varying amplification efficiencies and coverage rates of the primers. The bacteria in rare biospheres can never be identified if the primers used are not applicable to them.

Previous studies have shown that diverse microbial body structures can inhabit hot springs with a wide temperature range from ambient to boiling. (Barns et al., 1994; Pace 1997; Bonch-Osmolovskaya et al., 1999; Marteinsson et al., 2001; Kumar et al., 2004; Meyer-Dombard et al., 2005; Lau et al., 2006, 2009; Hall et al., 2008; Aditiawati et al., 2009; Mitchell 2009; Reigstad et al., 2009; Song et al., 2009, 2010; Aguilera et al., 2010; Jiang et al., 2010; Sayeh et al., 2010; Vick et al., 2010). *In silico* analysis of obtained results revealed that majority of the total partial 16S rRNA gene fragments belong to domain bacteria, where as only one sequence belonged to domain Archaea whose phylum identified was *Woesearchaeota*. The bacterial community structure detected in present study is given in figure 4. An evident phylum *Proteobacteria* was most abundant, constituting 36% and 40% of the total bacteria identified using V5-6 and V3-6 regions, respectively. Phylum *Proteobacteria* comprised mainly of *hydrogenophilus* with *pseudomonas*, *sphingomonas*, *rhodovibrio*, *nitrospira*, *serpens*, *klebsiella* and *Escherichia/Shigella* being present in small portion. The majority of *Hydrogenophilus* were represented by the order *Hydrogenophilales*. *Hydrogenophilales* generally includes anaerobic, rod-shaped, gram-negative, chemolithoautotrophic bacteria. Other phyla *Thermodesulfobacteria* (9.3%), *Firmicutes* (7.5%), *Deinococci-thermus* (7%), *Bacterioidates* (2.5%), *Aquificae* (2%) were assigned to V5-6 region amplified sequences while phyla *Firmicutes* (18%), *Bacterioidates* (6%), *Chloroflexi* (2%) were assigned to V3-6 region amplified sequences. A significant number of sequences, 30.5% and 20.5% from region V3-6 and V5-6 respectively could not match to existing database which were recorded as unclassified Bacteria. The most dominant phylotypes inhabiting Tattapani hot water spring were found to be *Proteobacteria*, *Firmicutes*, *Thermodesulfobacterium* and *Deinococcus-Thermus*, which are widely distributed in both terrestrial and aquatic environment. Apart from this other phylotypes of this spring were *Bacterioidates*, *Chloroflexi* and *Aquificae*. The phylotypes *Proteobacteria*, *Firmicutes*, *Thermodesulfobacterium*, *Deinococcus-Thermus*, *Bacterioidates*, *Chloroflexi* and *Aquificae* have been reported to inhabit Siloam hot water spring, Limpopo, South Africa (Tekere et al., 2011); Malaysian hot spring (Chan et al., 2015); hot spring on the Tebetian Plateau, China (Huang et al., 2011).

In total 11 distinct classes were identified out of which 7 classes were belonged to region V3-6 and all 11 classes to region V5-6. All the classes from region V3-6 were detected in 16S rRNA fragments from region V5-6. The most dominant class of the phylum *Proteobacteria* in this hot water spring was the *betaproteobacteria* (78%) followed by *gmmaproteobacteria* (19%) and *alphaproteobacteria* (4%). Other major classes detected for both regions were *Clostridia*, *Sphingobacteria*, *Bacilli*, *Anaerolineae*, *unclassified Bacteroidetes* and *unclassified Firmicutes*. Classes unique to region V5-6 were *Thermodesulfobacteria*, *Deinococci*, *Cytophagia* and *Aquificae*.

Out of 15 distinct identified orders, 12 orders belonged to V5-6 region and 9 orders were obtained from V3-6 region. The order *Hydrogenophilales* was found to be most abundant in both the regions. Other major orders identified for region V5-6 were *Thermodesulfobacteriales*, *Clostridiales*, *Pseudomonadales*, *Thermales*, *Enterobacteriales*, *Desulfurococcales*, *Pseudomonadales* and *Bacillales* and for region V3-6 were *Clostridiales*, *Thermoanaerobacterales*, *Sphingobacteriales* and *Coriobacteriales*.

A total of 17 and 14 different genera were assigned to region V5-6 and V3-6 respectively. The most abundant bacterial genus identified using both variable regions was *Hydrogenophilus*. Other detectable genera for region V5-6 were *Caldimicrobium*, *Thermus*, *Pseudomonas*, *Hyperthermus* etc., and for region V3-6 were *Fervidicola*, *Desulfoviregula*, *Salisaeta*, *Thermobrachium*, *Rhodothermus* etc. Although, the dominant microbial groups present in extreme environments are more emphasised, it is worth noting that the less abundant microbes found in extreme environment could be an important individuals of the unexplored community which can reuse as unculturable genetic resource (Sogin et al., 2006).

CONCLUSIONS

Analysis of DNA directly extracted from environmental samples allows us to study natural microbial communities without the need for cultivation. The spring water rich in evident bacterial phylum *Proteobacteria* and other bacterial phyla found in the present study were *Thermodesulfobacteria*, *Firmicutes*, *Deinococci-thermus*, *Bacteroidates* and *Aquificae*. Isolation of genes responsible for high temperature tolerance in these bacteria can be used for developing high temperature tolerance in crop plants. This study will act as a foundation for further studies on microbial community diversity of extreme habitats.

ACKNOWLEDGEMENTS

The authors thank to Dr. T. R. Sharma, Project Director ICAR-NRCPB, New Delhi. Special thanks to Dr. Vinay Sharma ICAR-CPRI, Shimla for their help during sample collection.

REFERENCES

1. Abou-Shanab, R. A. I. (2007). Characterization and 16S rDNA identification of thermo-tolerant bacteria isolated from hot springs. *J. Appl. Sci. Res.* 3: 994-1000
2. Aditiawati, P., Yohandini, H., Madayanti, F., & Akhmaloka. (2009). Microbial diversity of acidic hot spring (kawah hujan B) in geothermal field of kamojang area, west Java-Indonesia. *Open Microbiol J* 3:58-66
3. Aguilera, A., Souza-Egipsy, V., Gonzalez-Toril, E., Rendueles, O., & Amils, R. (2010). Eukaryotic microbial diversity of phototrophic microbial mats in two Icelandic geothermal hot springs. *Int Microbiol* 13:21-32
4. Amann, R. I., Ludwig, W., & Schleifer, K. H. (1995). Phylogenetic identification and in situ detection of individual microbial cells without cultivation. *Microbiol. Rev.* 59: 143-169
5. Armougom, F., & Raoult, D. (2009). Exploring microbial diversity using 16S rRNA high-throughput methods. *J. Compu. Sci. Syst. Biol.* 2: 74-92
6. Barns, S. M., Fundyga, R. E., Jeffries, M. W., & Pace, N. R. (1994). Remarkable archaeal diversity detected in a Yellowstone National Park hot spring environment. *Proc Natl Acad Sci USA* 91:1609-1613
7. Becker, S., Singh, A. K., Postius, C., Boger, P., & Ernst, A. (2004). Genetic diversity and distribution of periphytic *Synechococcus* spp. in biofilms and picoplankton of Lake Constance. *FEMS Microbiol. Ecol.* 49: 181-190

8. Bonch-Osmolovskaya, E. A., Miroshnichenko, M. L., Slobodkin, A. I., Sokolova T. G., Karpov, G. A., Kostrikina, N. A., Zavarzina, D. G., Prokof'eva, M. I., Rusanov, I. I., & Pimenov, N. V. (1999). Biodiversity of anaerobic lithotrophic prokaryotes in terrestrial hot springs of Kamchatka. *Microbiology* 68:343–351
9. Briggs, B. R., Brodie, E. L., Tom, L. M., Dong, H., Jiang, H., & Huang, Q., (2014). Seasonal patterns in microbial communities inhabiting the hot springs of Tengchong, Yunnan Province, China. *Environ. Microbiol.* 16(6):1579-91
10. Chan, C. S., Chan, K. G., Tay, Y. L., Chua, Y. H., & Goh, K. M. (2015). Diversity of thermophiles in a Malaysian hot spring determined using 16S rRNA and shotgun metagenome sequencing. *Front Microbiol.* 5: 6:177
11. Cole, J. K., Peacock, J. P., Dodsworth, J. A., Williams, A. J., Thompson, D. B., Dong, H., Wu, G., & Hedlund, P., B. (2013). Sediment microbial communities in Great Boiling Spring are controlled by temperature and distinct from water communities. *ISME J* 7: 718–729
12. Hall, J. R., Mitchell, K. R., Jackson-Weaver, O., Kooser, A. S., Cron, B. R., Crossey, L. J., & Takacs-Vesbach, C. D. (2008). Molecular characterization of the diversity and distribution of a thermal spring microbial community by using rRNA and metabolic genes. *Appl Environ Microbiol* 74:4910–4922
13. Hou, W., Wang, S., Dong, H., Jiang, H., Briggs, B. R., Peacock, J. P., Huang, Q., Huang, L., Wu, G., Zhi, X., Li, W., Dodsworth, A. J., Hedlund, P. B., Zhang, C., Hartnett, E. H., Dijkstra, P., & Hungate, A. B. (2013) A Comprehensive Census of Microbial Diversity in Hot Springs of Tengchong, Yunnan Province China Using 16S rRNA Gene Pyrosequencing. *PLoS ONE* 8(1): e53350.
14. Huang, Q., Dong, C. Z., Dong, R. M., Jiang, H., Wang, S., Wang, G., Fang, B., Ding, X., Niu, L., Li, X., Zhang, C., & Dong, H. (2011). Archaeal and bacterial diversity in hot springs on the Tibetan Plateau, China. *Extremophiles.* 15(5):549-63
15. Huber, J. A., Mark, Welch, D. B., Morrison, H. G., Huse, S. M., & Neal, P. R. (2007). Microbial population structures in the deep marine biosphere. *Science* 318: 97–100
16. Huse, S. M., Dethlefsen, L., Huber, J. A., Welch, D. M., Relman, D. A., & Sogin, M. L. (2008). Exploring microbial diversity and taxonomy using ssu rna hypervariable tag sequencing. *PLoS Genet.* 4: e1000255
17. Jiang, H., Huang, Q., Dong, H., Wang, P., Wang, F., Li, W., & Zhang, C. (2010). RNA-based investigation of ammonia-oxidizing archaea in hot springs of Yunnan Province, China. *Appl Environ Microbiol* 76:4538–4541
18. Kumar, B., Trivedi, P., Mishra, A. K., Pandey, A., & Palni, L. M. S. (2004). Microbial diversity of soil from two hot springs in Uttarakhand Himalaya. *Microbiol Res* 159:141–146
19. Lau, M. C. Y., Aitchison, J. C., & Pointing, S. B. (2009). Bacterial community composition in thermophilic microbial mats from five hot springs in central Tibet. *Extremophiles* 13:139–149
20. Lau, M. C. Y., Jing, H., Aitchison, J. C., & Pointing, S. B. (2006). Highly diverse community structure in a remote central Tibetan geothermal spring does not display monotonic variation to thermal stress. *FEMS Microbiol Ecol* 57:80–91
21. Liu, Z., DeSantis, T. Z., Andersen, G. L., & Knight, R. (2008). Accurate taxonomy assignments from 16S rRNA sequences produced by highly parallel pyrosequencers. *Nucleic Acids Res.* 36: 120
22. Mangrola, A., Dudhagara, P., Koringa, P., Joshi, C. G., Parmar, M., & Patel, R. (2015). Deciphering the microbiota of Tuwa hot spring, India using shotgun metagenomic sequencing approach. *Genomics data* 153-155
23. Marteinsson, V. T., Hauksdottir, S., Hobel, C. F. V., Kristmannsdottir, H., Hreggvidsson, G. O., & Kristjansson, J. K. (2001). Phylogenetic diversity analysis of subterranean hot springs in Iceland. *Appl Environ Microbiol* 67:4242–4248

24. Meyer-Dombard, D. R., Shock, E. L., & Amend, J. P. (2005). Archaeal and bacterial communities in geochemically diverse hot springs of Yellowstone National Park, USA. *Geobiology* 3:211–227
25. Mitchell, K. R., (2009). *Controls on microbial community structure in thermal environments: exploring bacterial diversity and the relative influence of geochemistry and geography*. University of New Mexico, Mexico.
26. Muyzer, G. E., Waal, C., & Uitterlinden, A. G. (1993). Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis and polymerase chain reaction-amplified genes coding for 16S rRNA. *Appl. Environ. Microbiol.* 59: 695–700
27. Narayan, V. V., Hatha, M. A., Morgan, H. W., & Rao, D. (2008). Isolation and characterisation of aerobic thermophilic bacteria from the savusavu hot spring in Fiji. *Microbes Environ.* 23: 350- 352
28. Naruya, S. & Masatoshi, N. (1987). *The Neighbor-joining Method: A New Method for Reconstructing Phylogenetic Trees*. *Mol. Biol. Evol.* 4(4):406-425
29. Nichols, D. S., Sanderson, K., Buia A., Kamp, J., Halloway, P., Bowman, J. P., Smith, M., Mancuso, C., Nichols, P. D., & McMeekin, T. A. (2002). Bioprospecting and biotechnology in Antarctica. In: Jabour G J, Haward M (eds). *The Antarctic: past, present and future*. Antarctic CRC. Res. Report Hobart. 28: 85-103
30. Pace, N. R. (1997). A molecular view of microbial diversity and the biosphere. *Science* 276:734–740
31. Reigstad, L. J., Jorgensen, S. L., & Schleper, C. (2009). Diversity and abundance of Korarchaeota in terrestrial hot springs of Iceland and Kamchatka. *ISME J* 4:346–356
32. Roesch, L. F. W., Fulthorpe, R. R., Riva, A., Casella, G., Hadwin, A. K. M., Kent, A. D., Daroub, S. H., Camargo, F. A. O., Farmerie, W. G., & Triplett, E. W. (2007). Pyrosequencing enumerates and contrasts soil microbial diversity. *Int. Soc. Microb. Ecol. J.* 1: 283-290
33. Sambrook and Russell (2001) *Molecular Cloning a Laboratory Manual*
34. Sayeh, R., Birrien, J. L., Alain, K., Barbier, G., Hamdi, M., & Prieur, D. (2010). Microbial diversity in Tunisian geothermal springs as detected by molecular and culture-based approaches. *Extremophiles* 14:501–514
35. Sharma, P. K., Caplash, N., & Kaur, J. (2007). An improved method for single step purification of metagenomic DNA. *Mol. Biotechnol.* 36: 61-63
36. Sogin, M. L., Morrison, H. G., Huber, J. A., Welch, D. M., & Huse, S. M. (2006). Microbial diversity in the deep sea and the underexplored rare biosphere. *Proc. Natl. Acad. Sci. USA* 103: 12115–12120
37. Song, Z. Q., Chen, J. Q., Jiang, H. C., Zhou, E. M, Tang, S. K, Zhi, X. Y., Zhang, L. X., Zhang, C. L. L., & Li, W. J. (2010). Diversity of Crenarchaeota in terrestrial hot springs in Tengchong, China. *Extremophiles* 14:287–296
38. Song, Z., Jiang, H., Zhi, X., Zhang, C., Dong, H., & Li, W. (2009). Actinobacterial diversity in hot springs in Tengchong (China), Kamchatka (Russia), and Nevada (USA). *Geomicrobiol J* 26:256–263
39. Tekere, M., Adele, L., Jana, O., Nelia, J., & Stephanus, V. (2011). Metagenomic analysis of bacterial diversity of Siloam hot water spring, Limpopo, South Africa. *Afr. J. Biotechnol.* 10: 18005-18012
40. Vick, T. J., Dodsworth, J. A., Costa, K. C., Shock, E. L., & Hedlund, B. P. (2010). Microbiology and geochemistry of Little Hot Creek, a hot spring environment in the Long Valley Caldera. *Geobiology* 8:140–154
41. Wang, Y., & Qian, P. (2009). Conservative fragments in bacterial 16S rRNA gene and primer design for 16S ribosomal DNA amplicons in metagenomic studies. *PLoS one* 4:10:e7401

42. Wang, Y., & Qian, P. Y. (2009). Conservative fragments in bacterial 16S rRNA genes and primer design for 16S rribosomal DNA amplicons in metagenomic studies. *PLoS One*. 4: 7401-7409
43. Wright, E. S., Yilmaz, L. S., & Noguera, D. R. (2012). DECIPHER, A Search-Based Approach to Chimera Identification for 16S rRNA Sequences. *Applied and Environmental Microbiology* 78(3):717-25
44. Zhou, J., Bruns, M. A., & Tiedje, J. M. (1996). DNA recovery from soils of diverse composition. *Appl. Enviro. Microbiol.* 62:316–322

APPENDICES



Figure 1: Location of Hot Water Spring Tattapani, Himachal Pradesh.
 (a)-Google Map Showing; Hot Water Location of the Spring; (b)-Actual Sites of the Sampling of Hot Water




Figure 2: Agarose Gel Electrophoresis a-(Lane 1-3) Metagenomic DNA and Lane M- Lambda DNA /EcoRI+HindIII marker. b- (Lane 1,2) Amplicon Obtained Using Primer Pair U789F-U1053R; (Lane 3,4) Amplicons Obtained Using Primer Pair MGUF-MGUR




Figure 3a: Neighbor-Joining Tree Showing the Phylogenetic Relationships of Partial 16S rRNA Gene Sequences Amplified from V3-V6 Region




Figure 3b: Neighbor-Joining Tree Showing the phylogenetic Relationships of Partial 16S rRNA Gene Sequences Amplified from V5-V6 Region




Figure 4: Bacterial Community Structure Obtained from V3-6 (a-d) and V5-6 (e-f) Region; a & e – Phyla, b & f- Classes, c & g –Order, d & h- Genera